

Prevalence of *Corynespora* leaf spot disease of cotton under South Gujarat of India

Nirva Patel^{1*}, Prashant B. Sandipan², Nishi Saini³, P. S. Patel² and R. K. Patel⁴

¹Department of Plant Pathology, N. M. College of Agriculture, Navsari Agricultural University (NAU), Navsari 396 450 (Gujarat), India

²Main Cotton Research Station (MCRS), Navsari Agricultural University (NAU), Surat 395 007 (Gujarat), India

³Department of Genetics and Plant Breeding, Maharana Pratap University of Agriculture and Technology (MPUAT), Udaipur 313 001 (Rajasthan), India

⁴Krishi Vigyan Kendra, Navsari Agricultural University (NAU), Surat 395 007 (Gujarat), India

Citation: Nirva Patel, Prashant B. Sandipan, Nishi Saini, P. S. Patel, and R. K. Patel (2023). Prevalence of *Corynespora* leaf spot disease of cotton under South Gujarat of India. *Acta Botanica Plantae* V02i02, 18-23. DOI: <http://dx.doi.org/10.5281/zenodo.8050114>

Corresponding Author: Nirva Patel | E-Mail: (pnirvapatel13@gmail.com)

Received 15 January 2023 | Revised 22 March 2023 | Accepted 28 April 2023 | Available Online April 27 2023

Copyright: This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Cotton is one of the most important economic products of the group of fibers due to volume and value of production. Its cultivation is also of great social importance, due to the number of jobs generated directly or indirectly. Cotton is an important fiber-yielding crop of global importance, which is grown in tropical and subtropical regions of more than 80 countries of the world. In this experiment, prevalence and incidence were recorded in the cotton growing regions of Surat, Bharuch, and Narmada districts of South Gujarat in the year 2021 to examine the presence of the *Corynesporacassiicola* pathogen on cotton plants and to record the observations on Per cent Disease Incidence (PDI). The maximum percent disease intensity was found in the Surat district, while the lowest disease intensity of *C. cassiicola* was found in the Bharuch district.

Keywords: Cotton, *Gossypium*, Prevalence, Incidence, *Corynespora*, Village, District, Intensity, Diseases

INTRODUCTION

Cotton (*Gossypium hirsutum* L.) is one of the most important fiber crops playing a key role in the economic and social scenario of the globe. It is the oldest among all the commercial crops of the world, providing fiber for the clothing of mankind. It is also known as "The white gold" or "The king of fibers". Cotton is a premier cash crop of our country and belongs to the family Malvaceae [1]. It is one of the most ancient and important commercial crops next only to food grains and is the principal raw material for flourishing the textile industry. It provides employment and sustenance to a population of nearly 42 million people, who are involved directly or indirectly in cotton production, processing, textiles and related activities [15]. Cotton is grown worldwide for its natural fiber and oil. The cotton seed contains 30 percent starch, 25 percent oil, and 16.20 percent protein. It is also being used in the manufacture of medicinal supplies, tarpaulin, cordage and belting. The cotton hulls serve as roughage for livestock and the fuzz (short seed hair) is used in the manufacture of papers, plastics, carpets, rayon, explosives and cotton wools etc. [18]. India is one of the major cotton growing countries in the world. India ranks first in area and second in the total production of cotton in the world. Hence, India has a large domestic textile

industry. It is chiefly grown in Maharashtra, Gujarat, Andhra Pradesh, Madhya Pradesh, Punjab, Tamil Nadu and Karnataka. India is the largest cotton-growing country in the world with an area of around 134.77 lakh hectares with production of 360.65 lakh bales and productivity of 455kg/ha [2]. In Gujarat, cotton is cultivated in an area of 26.55 lakh hectares and the production of 86.17 lakh bales with a productivity 552kg/ha [2].

In India, the productivity of cotton is very low due to many constraints including diseases. Diseases are inherent compounds of the agroecosystem that must be dealt with continuously and on a knowledge basis. Cotton is affected by various diseases caused by fungi, bacteria and viruses. The most common cotton diseases reported in India are Wilt (*F. oxysporum* f. sp. *vasinfectum* (G.F. Atk.) W.C. Snyder & H.N. Hansen), Root rot (*Rhizoctonia bataticola* (Taubenh.), *Verticillium* wilt (*Verticillium dahliae* Kleb.), Anthracnose (*Colletotrichum gossypii* Southworth or *C. capsici* (Syd.) Butler & Bisby), Gray mildew (*Ramularia areola* G.F. Atk.), Blackarm (*Xanthomonas campestris* sp. *malvacearum* (Pammel) Dowson), *Corynespora* leaf blight (*Corynespora cassiicola* (Berk. & M. A. Curtis) C. T. Wei, Leaf blight (*Alternaria*

macrosporaZimm) and Leaf curl (Cotton leaf curl virus), Boll rot [19] and Physiological disorders as Para wilt, Leaf reddening and sometimes Leaf elongation due to improper use of weedcides etc. Among them, Corynespora leaf spot of cotton is one of the important and serious diseases of cotton. Looking to the history target leaf spot caused by *C.cassicola* was reported in Alabama for the first time in 1959 on upland cotton (*Gossypium hirsutum* L.), the most widely planted species of cotton in the world [13 and [17]. In 1995, target spot was reported on upland cotton in Brazil but under the name of Corynespora leaf spot [16]. The first occurrence and re-emergence of the disease on upland cotton appeared in China [25], Brazil [12], and several states in the United States including Georgia [11], Alabama [8] and [9], China [25], Brazil [12] and [10].Louisiana [20], Tennessee [7], Central India [21] and South India [6]. Target spot has been a concern for farmers and researchers due to its increasing occurrence especially on cotton [23] owing to the monoculture farming, adoption of conservation tillage systems, susceptibility of current cultivars,

lack of crop rotation and optimal weather patterns for disease development [14] and [5].

The initial symptoms of target spots in cotton are characterized by small spots on the leaves located in the lower stratum of the plant [9]. The symptoms were observed in the lower canopy, which progressed upward to cover the entire plant. Initially, leaves exhibited circular to irregular, dark red, small and numerous lesions, which over the time became brown (5-10mm) surrounded by a dark border. As lesions matured, alternating rings of light and dark brown bands developed. The most mature lesions presented like a target-type appearance. Lesions may present as concentric rings [11] and in case of great severity, the leaves acquire a yellowish colour and easily detach from the branches resulting in defoliation [9]. Looking to the recurrent occurrence in one or other region and inflicting intensity or damage under the South Gujarat region. Hence, the present investigation was undertaken to illuminate deeply and meticulously this disease in these districts of South Gujarat.

Word cloud prepared from the Introduction part



Material and Methods

In the South Gujarat region, a survey of the major diseases of cotton crops was carried out in different fields in various districts. Cotton fields from different areas were selected randomly on the survey route. In each field, plants were selected at random and the severity of different diseases was recorded. The range of severity of disease intensity was calculated for foliar disease as the incidence of particular diseases in their respective areas.

Five leaves from the lower part and 5 leaves from the middle/ plant were selected by using 0-4 scale as given by [22] and then these grades were converted into percent disease intensity (PDI) by using the formula given by [26].

$$\text{Disease incidence (\%)} = \frac{\text{No. of infected plants}}{\text{No. of leaves observed} \times \text{Max. Grade}} \times 100$$

Disease Rating Scale for *Alternaria* (ALS), *Corynespora* (CoLS), *Myrothecium* (MLS) and Rust disease

Scale	PDI	Grade	Symptoms
0	0.0	Immune	No Infection
1	1-25%	R	A few small spots, less than 2mm, scattered, which over less than 5% leaf area
2	26-50%	MR	Spots bigger in size up to 3mm and cover 6-20% leaf area covered
3	51-75%	MS	Spots increasing in size 3-5mm, irregular in shape, coalescing and 21-40% leaf area cover
4	>75%	S	Many spots coalesce to make bigger lesion, irregular in shape and size and covering more than 40% leaf area

For foliar diseases, it is the standard methodology of AICRP, Cotton.

Table: 1 Information details for South Gujarat Farmers

1	Date of sowing	:	Farmer's field: With onset of monsoon Research Farm: 12 th -29 th June (Depending upon rain)
2	Normal or delayed sowing	:	-
3	If delayed, reasons there of	:	-
4	Number of entries sown	:	BG II hybrids on farmer's field/ others, if any
5	Details of additional entries sown, if any	:	-
6	Whether border rows sown	:	-
7	No. of rows/entry	:	-
8	Row length (in m)	:	-
9	Spacing (in cm)	:	120 x 45 (Irrigated) and 90 x 30 (Rainfed)
10	No. of replications	:	-
11	Irrigation No. with date	:	As per availability
12	No. of Weeding	:	Pre-emergence application of herbicide + 2-3 manual weeding + 2-3 interculturing
13	Fertilizer application	:	240-40-00 NPK kg/ha as per recommendation for hybrids, <i>G. hirsutum</i> 120-0-0 kg/ha and <i>G. herbaceum</i> & <i>G. arboreum</i> 80-0-0 kg/ha (Rainfed)
14	Insecticide/Pesticide application	:	Insecticides for sucking pest & Pink boll worm management and fungicides for foliar diseases

Result & Discussions

In the years 2021, a roving survey was conducted to determine the presence of *C. cassiicola* pathogen on the cotton plants and to record the observations on Per cent Disease Intensity (PDI) in the cotton-growing regions of Surat, Bharuch and Narmada districts of South Gujarat. Five areas in each field were chosen to determine the severity of the disease. A total of 25 plants were evaluated from each farmer's field. The severity of the disease condition was determined by using a 0-4 point disease assessment scale as mentioned. The survey was conducted for symptomatology, incidence, distribution, and spread of the disease along with the results are provided in the Table 2.

Among all the districts of South Gujarat surveyed, the *C. cassiicola* was more prevalent in Surat district. Various locations had different soil and climatic conditions and crops were at different growth stages during the field visit. The cotton crops were surveyed for the prevalence of *C. cassiicola* pathogen. Photo 1 Micrograph showing conidia and

conidiophore of *Corynespora cassiicola*. In the Surat district, a survey was carried out in three locations which, had medium to black soil type and the crop was in the flowering to fruiting stage. Table 2 and Fig. 1 revealed that during the year 2021 the disease intensity of Surat district was ranged from 1.50 to 26.50 per cent. Maximum disease intensity of 0.00-26.50 percent was observed in LRA 5166 followed by *G. Cot. Hy. 12 BG II* with 0.00-8.50 PDI, *G. Cot. Hy. 8 BG II* with 0.00-6.50 PDI and *G. Cot. Hy. 10 BG II* with 0.00-5.50 PDI at Main Cotton Research Station (MCRS) of Choryasi taluka of Surat. The average disease intensity of 17.00 percent was observed in Surat district. In Bharuch district, total of three locations were surveyed. The location had black soil and the cotton crop was in the flowering to the fruiting stage and the disease intensity was ranged from 1.50 to 3.50 percent with the maximum disease intensity of 0.00-3.50 percent was observed in Trinetra BGII at Chiklota village of Valia taluka and the average district disease intensity was 3.00 percent. In Narmada district, total three locations

were surveyed. The location had medium to black soil and cotton crop was in the flowering to fruiting stage and the disease intensity ranged from 1.00 to 6.50 percent. Maximum disease intensity of 0.00-6.50 per cent was observed in Sadabahar BG II at Kukarada village of Dediypada taluka and the average district disease intensity was 3.16 per cent. Overall, the survey results indicated that the MCRS location of Surat district, which can be considered as the main region for the presence of *C. cassiicola* pathogen.

Similar research findings were reported by [24], who carried out a survey of the target leaf spot disease in chilli in major chilli growing areas at Basantpur, Mohanpur, Haringhata, Jajuli, Kalyani and Laupala in Nadia district of West Bengal during the year 2015-16. The disease incidence range varied from 11 to 35 percent in different chilli fields.

The survey was conducted in the major cotton-growing districts of Guntur, Krishna, and Prakasam of Andhra Pradesh. The intensity of *Corynespora* leaf spot disease was observed 15 and 12 percent in Akira and RCH 659 in Krishna district. The average highest PDI 21.3 percent, 11.07 percent and 7.7 percent of *Corynespora* leaf spot was recorded in Krishna, Guntur and Prakasam districts [3]. The survey was conducted in major cotton-growing districts of Bhavnagar, Rajkot and Junagadh of Gujarat, 2021. In mid-season, the PDI of *Corynespora* leaf spot (CoLS) was observed in the range of 0.0-10.5 percent. Whereas, in the late season, PDI of the *Corynespora* leaf spot disease (CoLS) was recorded in the range of 15.0-55.2 percent [4]. Whereas, in the late season, PDI of the *Corynespora* leaf spot disease (CoLS) was recorded in the range of 15.0-55.2 percent [4].



(1) MCRS, Surat



(2) Modaliya, Bharuch



(3) Dediypada, Narmada



(4) Sagbara, Narmada

Table:2 Survey for occurrence of Target spot disease in cotton crop of South Gujarat during the year 2021.

Sr. No.	District	Taluka	Village / GPS Coordinates	Type of soil	Variety	Stage of crop	Percent Disease Intensity (PDI) Range
1.	Surat	Mangrol	Ambavadi 21.08°56" N, 72.71°71" E	Medium black	Pandav BG II	Flowering	0.00-2.50%
		Choryasi	MCRS farm 21.17°00" N, 72.80°10" E	Black	G. Cot. Hy. 12 BG II G. Cot. Hy. 8 BG II G. Cot. Hy. 10 BG II LRA 5166	Fruiting	0.00-8.50% 0.00-6.50% 0.00-5.50% 0.00-26.50%
		Umarpada	Vakrant Amba 21.46°56" N, 73.56°53" E	Medium black	Rashi Neo 578 BG II	Flowering	0.00-1.50%
	District Mean						
2.	Bharuch	Valia	Chiklota 21.63°38" N, 73.31°46" E	Black	Trinetra BG II	Fruiting	0.00-3.50%
		Netrang	Modaliya 21.21°55" N, 71.54°27" E	Black	Pandav BG II	Flowering	0.00-2.00%
		Amod	Ranipura 22.01°53" N, 72.87°81" E	Black	Shanar BG II	Fruiting	0.00-1.50%
	District Mean						
3.	Narmada	Rajpipala	Mota Limbatwada 21.87°57" N, 73.55°94" E	Medium Black	Bhagya 999 BG II	Flow ering	0.00-1.00%
		Dediypada	Kukarada 21.94°51" N, 73.87°42" E	Medium Black	Sadabahar BG II	Flowering	0.00-6.50%
		Sagbara	Simamli 21.52°00" N,	Black	Rashi 658 BG II	Flowering	0.00-2.00%




Fig. 1 Survey for target leafspot disease of cotton in South Gujarat during the year 2021

Acknowledgement

The author is highly thankful to the Department of Plant Pathology, N. M. College of Agriculture, NAU, Navsari (Gujarat) and the Scientists of Main Cotton Research Station (MCRS), NAU, Surat (Gujarat) for providing the requisite facility and other obligatory arrangements for the facilitating the experiment.

References

- Anonymous (2017a). <http://en.wikipedia.org/wiki/Cotton>.
- Anonymous (2017b). <https://www.worldatlas.com/articles/top-cotton-producing-countries-in-the-world.html>.
- Anonymous (2020b). All India Coordinated Research Project on Cotton – Annual Report., F-8.
- Anonymous (2021a) All India Coordinated Research Project on Cotton – Annual Report., F-9.
- Avozani, A.; Reis, E. M. and Tonin, R. B. (2014). Sensitivity loss by *Corynespora cassiicola* isolated from soybean to the fungicide carbendazim. *Summa Phytopathol.*, 40(3): 273-276.
- Bandi, M. V. S. P.; Bhattiprolu, S. L.; Kumari, V. P.; Kumar, V. M.; Divyamani, V.; Patibanda, A. K.; Jayalalitha, K. and Kumar, D. V. S. R. (2022). First report of *Corynespora cassiicola* causing target spot on cotton (*Gossypium hirsutum*) in South India. *Plant Dis.*, 106(8). DOI: 10.1094/pdis-11-21-2597-pdn.
- Butler, S.; Young-Kelly, H.; Raper, T.; Cochran, A.; Jordan, J.; Shrestha, S.; Lamour K.; Mengistu, A.; Castro-Rocha, A. and Shelby, P. (2016). First report of target spot caused by *Corynespora cassiicola* on cotton in Tennessee, USA. *J. Phytopathol. and Plant. Prot.*, 50(1): 18-29.
- Campbell, H. L.; Hagan, A. K.; Bowen, K. L. and Nightengale, S. P. (2012). *Corynespora* leaf spot: a new disease in Alabama cotton. *Phytopathol.*, 102(7): 18-19.
- Conner, K. N.; Hagan, A. K. and Zhang, L. (2013). First report of *Corynespora cassiicola* in Alabama. *Plant Dis.*, 97(10): 1379.
- Dias, A. R.; Souza, H. M. and Theodoro, G. F. (2016). Alvo mancha alvo em algodao. *Cult. Grand Cultur.*, 17: 18-21.
- Fulmer, A. M.; Walls, J. T.; Dutta, B.; Parkunan, V.; Brock, J. and Kemerait, J. C. (2012). First report of target spot caused by *Corynespora cassiicola* on cotton in Georgia. *Plant. Dis.*, 96(7): 1066-1066.
- Galbieri, R.; Araujo, D. C. E. B.; Kobayashi, L.; Giroto, L.; Matos, J. N.; Marangoni, M. S.; Almeida, W. P. and Mehta, Y. R. (2014). *Corynespora* leaf blight of cotton in Brazil and its management. *Americ. J. Plant Sci.*, 5(26): 3805-3811.
- Jones, J. P. (1961). A leaf spot of cotton caused by *Corynespora Cassiicola*. *Phytopathol.*, 51(5): 305-308.
- Koenning, S. R.; Creswell, T. C.; Dunphy, E. J.; Sikora, E. J. and Mueller, J. D. (2006). Increased occurrence of target spot of soybean caused by *Corynespora cassiicola* in the Southeastern United States. *Plant Dis.*, 90(7): 974.
- Manickam, S. and Sankaranarayana, K. (2013). Cotton Technical Assistance Programme for Africa. Training Manual on Relevance and Techniques of Organic Cotton Production. pp-1.
- Mehta, Y. R.; Motomura, K. F. and Almeida, W. P. (2005). *Corynespora* leaf spot of cotton in Fitopatologia Brasileira, 30: 131.
- Onesirosan, P. T.; Mabuni, C. T.; Durbin, R. D.; Morin, R. B.; Rich, D. H. and Arny, D. C. (1975). Toxin production by *Corynespora cassiicola*. *Physiol. Plant. Pathol.*, 5(3): 289-295.

18. Prasad, J. (2015). Studies on root rot of cotton (*Gossypium arboreum* L.) incited by *Macrophomina phaseolina* (Tassi) Gold.) with special reference to rhizosphere relationship. Thesis M. Sc. (Agri.), Rajasthan Agricultural University, Bikaner.
19. Prashant B. Sandipan, P. S. Patel, R. K. Patel, Rameela I. Chaudhari and M. C. Patel. 2022. Evaluation of different fungicides against boll rot and foliar diseases of cotton under South Gujarat of India. *Journal of Plant Development Sciences*, 14 (5): 491-496.
20. Price, P.; Singh, R. and Fromme, D. (2015). First report of target spot caused by *Corynespora cassiicola* in Louisiana cotton. *Plant Health Prog.*, 16(4): 223-224.
21. Salunkhe, V. N.; Gawande, S. P.; Nagrale, T.; Hiremani, N.S.; Gokte-Narkhedkar, N. and aghmare, V. N. (2019). First report of *Corynespora* leaf spot of cotton caused by *Corynespora cassiicola* in Central India. *Plant Dis.*, 103(7): 1785-1787.
22. Sheoraj. (1989). Grading system for cotton disease. CICR publication, Nagpur.
23. Sumabat, L. G.; Kemerait, R. C. and Brewer, M. T. (2018). Phylogenetic diversity and host specialization of *Corynespora cassiicola* responsible for emerging target spot disease of cotton and other crops in the Southeastern United States. *Phytopathol.*, 108(7): 892-901.
24. Suresh, V.; Sumalatha, N.; Ravat, V. K. and Basu, A. (2017). First report of Louisiana State Univ. Agric. Cent. Internet Bengal, India. *Int. J. Curr. Microbiol. App. Sci.*, 6(8): 3216-3219.
25. Wei, Y. X.; Zhang, H.; Pu, L. and Lui, X. M. (2014). First report of target spot of cotton caused by *Corynespora cassiicola* in China. *Plant Dis.*, 98(7): 1006.
26. Wheeler, B. E. J. 1969. *An Introduction to Plant Disease*. John Willey and Sons, London, pages: 374.